

USDA United States
Department of
Agriculture

Forest Service

Northeastern
Research Station

General Technical
Report NE-316



Proceedings

14th Central Hardwood Forest Conference

March 16-19, 2004
Wooster, Ohio



14th

CENTRAL HARDWOOD FOREST CONFERENCE

NOTICE:

File size is 9.5MB, 539 pages

To access the Table of Contents linked to individual
papers suitable for reprints, go to:

[http://www.fs.fed.us/ne/newtown_square/publications/
technical_reports/pdfs/2004/316papers/gtr316toc.pdf](http://www.fs.fed.us/ne/newtown_square/publications/technical_reports/pdfs/2004/316papers/gtr316toc.pdf)

ECOSYSTEM RESTORATION AND WILDFIRE MANAGEMENT TREATMENTS AFFECT SOIL ORGANIC MATTER AND MICROBIAL ACTIVITY IN FOUR CONTRASTING FORESTS

*R.E.J. Boerner, Thomas A. Waldrop, Carl N. Skinner, Mac A. Callabam, Jr.,
Jennifer A. Brinkman, and Annemarie Smith†*

As part of a national-scale evaluation of the consequences of restoration and wildfire fuel reduction treatments in ecosystems that historically had frequent fire (www.ffs.fs.fed.us), we determined the effects of reintroduction of dormant season fire (functional restoration) and thinning from below (structural restoration) on soil organic matter characteristics and microbial activity in two mixed oak forests in the central hardwoods region, and contrasted them with the effects of the same treatments in two conifer forests. The hardwood forests were located on the Allegheny Plateau of southern Ohio (Ohio Hills/OH) and in the southern Appalachian Mountains of North Carolina (Green River/NC). The conifer forests were a pine-oak site in the piedmont of South Carolina (Clemson Forest/SC) and a pine-fir site in the southern Cascades of northern California (Goosenest Adaptive Management Area/CA).

At OH and SC fire resulted in reduced soil organic C and total soil N, both alone and in combination with thinning. In contrast, thinning alone did not affect either C or N at OH, SC, or CA. Soil C:N ratio decreased in the plots that were burned at OH and SC, but not in those that were thinned or thinned+burned; at CA soil C:N ratio decreased in response to thinning. Fire, thinning, and the combination of the two resulted in reductions in acid phosphatase activity in soils at OH and SC, whereas at CA only the combination of fire and thinning reduced this metric of overall microbial activity. Chitinase activity was reduced by all three treatments at OH and SC. At CA, chitinase activity decreased in the order Control>Thin>Thin+Burn. Phenol oxidase activity changed little, and did not differ among treatment types at OH, SC, or CA.

Multiple regression models predict that posttreatment enzyme activity in these forests is more closely linked to changes in organic matter quality (soil C storage) and organic matter quality (N content, C:N ratio, phosphatase:phenol oxidase ratio) produced by the experimental treatments than to macroclimate, landscape-scale microclimate, geomorphological, or soil textural variations among treatment units within a site. These results suggest that the short term (1-2 yr) consequences of restoration treatments on soil microbial activity vary within and among ecosystem types in relation to their effects on the quality and quantity of soil organic matter, more so than in relation to vegetation, macroclimate, or geological setting.

This is contribution number 41 of the National Fire and Fire Surrogate Research Project. This research was funded through the National Fire Plan of the U.S.D.A. Forest Service. Although the authors received no direct funding from the U.S. Joint Fire Science Program (JFSP), this research could not have been accomplished without JFSP support of the existing FFS research project.

†Professor (REJB), Research Assistant II (JAB), Research Assistant I (AS), Department of Evolution, Ecology, and Organismal Biology, The Ohio State University, Columbus, OH 43210. Research Forester (TAW), U.S.D.A. Forest Service Southern Research Station, Clemson, SC 29634. Research Geographer (CNS), U.S.D.A. Forest Service Pacific Southwest Research Station, Redding, CA 96001. Research Ecologist (MAC), U.S.D.A. Forest Service Southern Research Station, Athens, GA 30602. REJB is corresponding author: to contact call (614) 292-6169 or e-mail at boerner.1@osu.edu.